

تقدم لجنة EiCoM الأكاديمية

حلول تقارير مختبر:

فيزياء عامة 1

جزيل الشكر للمهندس:

بشار العقابنة

LAB REPORT FOR EXPERIMENT 1

Date: 9/2/2017

Name: بشار عبد القادر العقاب

Partner's Name: _____

Registration No: 1440224

Registration No: _____

Physics Section: 16

Instructor's Name: سماء النمر

PHYSICS LAB EXPERIMENT 1: COLLECTION AND ANALYSIS OF DATA

1. PURPOSE :

(1)
We make this Imaginary Experiment to: know the relationship between the diameter of a hole in the bottom of the container [d], the depth of water in the container [h] and the time needed to empty the container [t].

II. DATA : (2) Convert Non-Linear relationship to Linear.

Table (1.1)

h (cm)	t in seconds			
	d = 1.5 mm	d = 2.0 mm	d = 3.0 mm	d = 5.0 mm
30.0	73.0	41.2	18.4	6.8
10.0	43.5	23.7	10.5	3.9
4.0	26.7	15.0	6.8	2.2
1.0	13.5	7.2	3.7	1.5

Using data in Table (1.1) fill in Table (1.2) below:

Table (1.2)

d (mm)	t in seconds			
	h = 30.0 cm	h = 10.0 cm	h = 4.0 cm	h = 1.0 cm
5.0	6.8	3.9	2.2	1.5
3.0	18.4	10.5	6.8	3.7
2.0	41.2	23.7	15	7.2
1.5	73.0	43.5	26.7	13.5

for $h = 30$ cm, fill in Table 1.3 below:

Table(1.3)

t (s)	d (mm)	1/d ² (mm ⁻²)
3.9	5.0	0.04
10.5	3.0	0.11
23.7	2.0	0.25
43.5	1.5	0.44

for $d = 2$ mm fill in Table 1.4 below:

Table(1.4)

log t	log h
0.568	0
0.833	0.6
1.021	1
1.265	1.5

III. ANALYSIS OF DATA :

Graph your results. **Independent** variables will be the diameter of hole and depth of water in the container. Time is the **dependent** variable and will depend on the previous two independent variables.


A. Plot the time (t) versus the depth (h) for each diameter (d) used. Do four graphs on one sheet, using the same set of axes, connecting points in a smooth curve for each and labeling them d_1 , d_2 , d_3 and d_4 .

B. On a second sheet of graph paper, plot the time (t) versus diameter (d) for each value of depth (h). Connect the points in a smooth curve and label the curves h_1 , h_2 , h_3 and h_4 .


C. Plot t versus 1/d² for $h = 30$ cm

D. plot log t versus log h for $d = 2$ mm.


A. The Relation between t and h at constant d [Table 1.1]




B. The Relation between t and d at constant h [Table I.2]



C. t versus $(1/d^2)$ for $h = 10\text{ cm}$ [Table 1.3]



D. $(\log(t))$ versus $(\log(h))$ for $d=3\text{ mm}$ [Table 1.4]



IV. CONCLUSIONS

1. From your graph (t) versus (h) for $d = 1.5$ mm, extrapolate the curve toward the origin. Does it pass through it? Would you expect it to do so?

Yes; because when there is no water in the container ($h=0$), the time needed to empty the container equals zero ($t=0$).

2. What type of relationship do you see between the time and diameter? Is it direct or inverse?

It's Inverse and non-Linear Relation.

3. From t versus $1/d^2$ graph, find the empirical relationship between time (t) and hole diameter (d) for $h = 30$ cm.

Slope (m) = $\frac{\Delta t}{\Delta (1/d^2)} = \frac{9.6 - 0}{0.1 - 0} = 96 \text{ s} \cdot \text{mm}^2$, y-intercept = 0

$\Rightarrow t = m \times \frac{1}{d^2} + \text{y-intercept} \Rightarrow t = 96 \times \frac{1}{d^2}$

4. From the previous relation, can you predict the time needed to empty the container if the diameter of the opening was 4 mm, 8 mm?

Yes; $t(4) = 96 (1/4^2) = 6 \text{ s}$

$t(8) = 96 (1/8^2) = 1.5 \text{ s}$

5. From the $\log t$ versus $\log h$ graph, find the empirical relationship between time (t) and depth (h) for $d = 2.3$ mm.

Slope (m) = $\frac{\Delta \log t}{\Delta \log h} = \frac{0.65 - 0.56}{0.2 - 0} = 0.45$, y-intercept = 0.56

$\Rightarrow \log t = 0.45 \log h + 0.56$

$\Rightarrow \log t - \log h^{0.45} = 0.56$

$\log (t/h^{0.45}) = 0.56$

$t/h^{0.45} = 10^{0.56} \Rightarrow t = 3.63 h^{0.45}$

find the slope on the graph

-0.5

6. Can you predict the time needed to empty the container if the depth of water was 25 cm, 80 cm?

Yes; $t(25) = 3.63 * (25)^{0.45} = 15.5 \text{ s}$

$t(80) = 3.63 * (80)^{0.45} = 26.1 \text{ s}$

LAB REPORT FOR EXPERIMENT 2

Date: 23/2/2017

Name: إسماعيل عبد القادر الكفاح

Partner's Name: _____

Registration No: 1440224

Registration No: _____

Physics Section: 16

Instructor's Name: سماء البشير

PHYSICS LAB EXPERIMENT 2 : MEASUREMENTS AND UNCERTAINTIES

I. PURPOSE :

- a) We will make some elementary measurement of length and mass, and from our measurements we will derive other quantities, such that: (volume, density,...)
- b) To estimate the inaccuracy or the error in each measurement.

II. DATA AND DATA ANALYSIS :

A. Measurement of π

Record your data in Table (2.1) below:

Table (2.1)

Trial No	By Caliber	$ d - \bar{d} $ (cm)	By a Piece of paper tape (نظم القوس)	$ c - \bar{c} $ (cm)
	d (cm)		c (cm)	
1	3.715	2.5×10^{-3}	12.0	0.1
2	3.710	2.5×10^{-3}	11.8	0.1
3	_____	_____	_____	_____
4	_____	_____	_____	_____
5	_____	_____	_____	_____
Average	$\bar{d} = 3.7125$ cm		$\bar{c} = 11.9$ cm	
Error	$\Delta \bar{d} = \pm 2.5 \times 10^{-3}$ cm		$\Delta \bar{c} = \pm 0.1$ cm	

1. Calculate the error, $\Delta \bar{d}$, in measuring the diameter of the disk, and, $\Delta \bar{c}$, in measuring the circumference and enter the values calculated in Table 2.1
Example for one calculation ($\Delta \bar{d}$) or ($\Delta \bar{c}$):

$$\Delta \bar{d} = \pm \sqrt{\frac{\sum_{i=1}^n |d_i - \bar{d}|^2}{n(n-1)}} = \sqrt{\frac{2 * (2.5 * 10^{-3})^2}{2(1)}} = 2.5 * 10^{-3} \text{ cm}$$

$$\Delta \bar{c} = \pm \sqrt{\frac{\sum_{i=1}^n |c_i - \bar{c}|^2}{n(n-1)}} = \sqrt{\frac{2 * (0.1)^2}{2(1)}} = 0.1 \text{ cm}$$

2. Using your average measured values of $\bar{d}$ and $\bar{c}$, calculate $\bar{\pi}$.

$$\bar{\pi}_{\text{exp}} = \frac{\bar{c}}{\bar{d}} = \frac{11.9}{3.7125} = 3.205$$

($c = \pi d$)

3. Calculate the error, $\Delta \bar{\pi}$, in the measured value, $\bar{\pi}$.

$$\text{Note: } \Delta \bar{\pi} = \bar{\pi} [(\Delta \bar{d} / \bar{d})^2 + (\Delta \bar{c} / \bar{c})^2]^{1/2} \quad (2.1)$$

$$\Delta \bar{\pi} = 3.205 (4.535 * 10^{-7} + 706.165 * 10^{-7})^{1/2} = 0.027$$

$$* \frac{\Delta \bar{d}}{\bar{d}} = 6.734 * 10^{-4} \quad * \frac{\Delta \bar{c}}{\bar{c}} = 8.403 * 10^{-3}$$

4. Which error contributes most to $\bar{\pi}$? (give a quantitative answer)

$$(\Delta \bar{d} / \bar{d} = 6.734 * 10^{-4}) < (\Delta \bar{c} / \bar{c} = 8.403 * 10^{-3}); \text{ as you can}$$

see the $(\Delta \bar{c} / \bar{c})$ contributes more than $(\Delta \bar{d} / \bar{d})$ to $(\bar{\pi})$, since it's the largest.

5. Does the measured average value of $\bar{\pi}$ agree with the accepted value of π (3.14159) within the calculated experimental error.

$$p.e. = \left| \frac{\bar{\pi}_{\text{exp}} - \pi_{\text{acc}}}{\pi_{\text{acc}}} \right| * 100 = \left| \frac{3.205 - 3.14159}{3.14159} \right| * 100\%$$

= 2.02% ; it's agree with the accepted value of π approximately.

B. Determination of Density

Record your data in Table (2.2) below:

Table (2.2)

Trial No	h (cm)	h - $\bar{h}$ (cm)	d (cm)	d - $\bar{d}$ (cm)
1	9.97	—	0.541	17.3×10^{-3}
2	—	—	0.500	23.7×10^{-3}
3	—	—	0.530	6.3×10^{-3}
4	—	—	—	—
5	—	—	—	—
Average	$\bar{h} = 9.97$ cm		$\bar{d} = 0.5237$ cm	
Error	$\Delta \bar{h} = \pm 2.5 \times 10^{-3}$ cm		$\Delta \bar{d} = \pm 12.252 \times 10^{-3}$ cm	
mass	$m = 16.15$ g		$\Delta m = \pm 0.01$ g	

1. Calculate the error, $\Delta \bar{h}$, in the average measured length and enter the result in Table (2.2).

$$\Delta \bar{h} = \frac{1}{2} (0.05) = 0.025 \text{ mm} = 2.5 \times 10^{-3} \text{ cm}$$

2. Calculate the error, $\Delta \bar{d}$, in the average measured diameter and enter the result in Table (2.2).

$$\Delta \bar{d} = \sqrt{\frac{\sum_{i=1}^n |d_i - \bar{d}|^2}{n(n-1)}} = \sqrt{\frac{9.0067 \times 10^{-4}}{6}} = 12.252 \times 10^{-3} \text{ cm}$$

خطأ لمتوسط التوزيع النتيجة

0.5

$\sqrt{(17.3 \times 10^{-3})^2 + (23.7 \times 10^{-3})^2 + (6.3 \times 10^{-3})^2}$

6

3. Take Δm to be half the smallest division of the balance used.
4. Using your average measured values of $\bar{h}$, $\bar{d}$, π determined in part A, and the measured value of mass m , calculate $\bar{\rho}$.

$$\bar{\rho} = \frac{4m}{\pi \bar{d}^2 \bar{h}} = \frac{4 \times 16.15}{3.14159 \times (0.5237)^2 \times 9.97} = 7.37 \text{ g/cm}^3$$

5. Calculate the error, $\Delta \bar{\rho}$, in the average value for the measured density, $\bar{\rho}$.

Note:
$$\Delta \bar{\rho} = \bar{\rho} \left[\left(\frac{\Delta m}{m} \right)^2 + \left(\frac{\Delta h}{h} \right)^2 + \left(\frac{2\Delta d}{d} \right)^2 + \left(\frac{\Delta \pi}{\pi} \right)^2 \right]^{1/2} \quad (2.2)$$

and use for $\bar{\pi}$ and $\Delta \pi$, the values determined in part A.

$$\Delta \bar{\rho} = 7.37 (3.834 \times 10^{-7} + 6.2876 \times 10^{-8} + 2.189 \times 10^{-3} + 7.097 \times 10^{-5})^{1/2}$$

$$= 0.35 \text{ g/cm}^3$$

6. Which error in m , $\bar{h}$, $\bar{d}$, or $\bar{\pi}$ contributes most to $\bar{\rho}$?
(give a quantitative answer)

$$\begin{aligned} * \frac{\Delta m}{m} &= 6.19195 \times 10^{-4} & * \frac{\Delta \bar{h}}{\bar{h}} &= 2.5075 \times 10^{-4} \\ * 2 \frac{\Delta \bar{d}}{\bar{d}} &= 46.79 \times 10^{-3} & * \frac{\Delta \bar{\pi}}{\bar{\pi}} &= 8.4243 \times 10^{-3} \end{aligned}$$

as you can see the $(2 \frac{\Delta \bar{d}}{\bar{d}})$ contributes most to $\bar{\rho}$, since it's the largest.

7. Using your calculations in (6), which error in m , $\bar{h}$, $\bar{d}$, or $\bar{\pi}$ contributes the least to $\bar{\rho}$?

$(\frac{\Delta \bar{h}}{\bar{h}})$ contributes the least to $\bar{\rho}$, because it's the smallest

8. Compare the measured value of $\bar{\rho} \pm \Delta \bar{\rho}$ with the accepted value of ρ ($\rho_{\text{accepted}} = 8.6 \text{ g/cm}^3$)

$$\text{p.e.} = \left| \frac{\bar{\rho}_{\text{exp}} - \rho_{\text{acc}}}{\rho_{\text{acc}}} \right| \times 100\% = \left| \frac{7.37 - 8.6}{8.6} \right| \times 100\%$$

$$= 14.3\%$$

LAB REPORT FOR EXPERIMENT 3

Date: 16/2/2017

Name: نسار عبد القادر الحفایه

Partner's Name: _____

Registration No: 1440224

Registration No: _____

Physics Section: 16

Instructor's Name: سماح نصر

PHYSICS LAB EXPERIMENT 3: VECTORS (FORCE TABLE)

I. PURPOSE :

* عمل (addition) لمجموعة من المتجهات بأكثر من طريقة

Measure the magnitude and direction of the resultant of three forces

experimentally by (the force table), graphically, and by the method of components.

II. DATA AND DATA ANALYSIS :

- ✗ Record the experimentally measured value of the resultant of the two forces in step one of the procedure (magnitude and direction) .

- ✗ Determine the resultant of the two forces in step one of the procedure graphically. How does it compare with the measured value?

- ✗ Again determine the resultant of the two forces in step one by the method of components. How does it compare with the measured value?

4. Record the experimentally measured value of the resultant of the three forces in step two of the procedure (magnitude and direction).

$$F_1 = 0.1 \times 9.8 = 0.98 \text{ N}, F_2 = 0.15 \times 9.8 = 1.47 \text{ N}, F_3 = 0.2 \times 9.8 = 1.96 \text{ N}$$

$$\theta_1 = 30^\circ \quad \theta_2 = 315^\circ \quad \theta_3 = 210^\circ$$

$$\Rightarrow F_R = 0.153 \times 9.8 = 1.4994 \text{ N}, \theta_R = 100 + 180 = 280^\circ$$

θ_{equ}

5. Determine the resultant of the three forces in step two of the procedure graphically using the polygon method. Compare it with the measured value.

Scale: 5 cm = 1 N, $F_{R \text{ graph}} = \frac{7.7 \text{ cm} \times 1 \text{ N}}{5 \text{ cm}} = 1.54 \text{ N}, \theta_R = 360 - 80.3 = 279.7^\circ$

$$\text{P.e} = \left| \frac{(F_{R \text{ graph}}) - (F_{R \text{ exp}})}{(1.54 + 1.4994) / 2} \right| \times 100\% = 2.67\%$$

* في المصكبة أن نقيس الزوايا

6. Again, use the method of components to determine the resultant for the three forces in step two of the procedure. Compare with experimental findings.

$$1^* F_1 = 0.98 \cos 30^\circ \hat{i} + 0.98 \sin 30^\circ \hat{j} = [0.85 \hat{i} + 0.49 \hat{j}] \text{ N}$$


$$F_2 = 1.47 \cos 315^\circ \hat{i} + 1.47 \sin 315^\circ \hat{j} = [1.04 \hat{i} + -1.04 \hat{j}] \text{ N}$$

$$F_3 = 1.96 \cos 210^\circ \hat{i} + 1.96 \sin 210^\circ \hat{j} = [-1.697 \hat{i} + -0.98 \hat{j}] \text{ N}$$

$$F_{Rx} = 0.85 + 1.04 - 1.697 = 0.193 \text{ N} \quad \rightarrow$$

$$F_{Ry} = 0.49 - 1.04 - 0.98 = -1.53 \text{ N} \quad \downarrow$$

$$\Rightarrow |F_R| = \sqrt{(0.193)^2 + (-1.53)^2} = 1.54 \text{ N}, \tan^{-1} \frac{1.53}{0.193} = 82.81^\circ \Rightarrow \theta = 277.19^\circ$$




$$2^* \text{P.e} = \left| \frac{1.54 - 1.4994}{(1.54 + 1.4994) / 2} \right| \times 100\% = 2.67\%$$

7. State the major source(s) of inaccuracy in the experimental results?

1. Friction between the string and the pulley.
2. Errors in reading the scale.
3. The force table may be not horizontal.

$5\text{cm} = 1\text{N}$



Kinematics of Rectilinear Motion

Name: نصار عبد القادر الحفالة

Instructor: سما 2 النمر

ID#: 1440224

Section: 16

Purpose: To study regular and irregular motion of objects by determining how the distance, velocity and acceleration are changing with time.

A. Fill in table (1) with data from your tape. Then, draw a graph of x versus t .

B. Use data from table (1) to fill table (2). Then, draw a graph of $\bar{v}$ versus t .

Draw the curve of the instantaneous velocity. {Assume that the instantaneous velocity at the interval's mid-point equals the average velocity in that interval, and that the acceleration is constant between each two interval.}

C. Use data from table (1) to fill table (3).

Table(1)

Total time t (sec)	Total distance x (cm)
0.0	0.0
0.1	2.7
0.2	6.1
0.3	9.6
0.4	13.3
0.5	17.1
0.6	20.9
0.7	24.2

Table(2)

«أخذ فترات متتالية» Time interval (s)	Average velocity $\bar{v} = \frac{\Delta x}{\Delta t}$ (cm/s)
0.0 - 0.1	27
0.1 - 0.2	34
0.2 - 0.3	35
0.3 - 0.4	37
0.4 - 0.5	38
0.5 - 0.6	38
0.6 - 0.7	33

Table (3)

t_{mid} (s)	$v_{mid} \cong \bar{v}$ (cm/s)	$\bar{a} \cong \frac{\Delta v}{\Delta t}$ (cm/s ²)
0.05	27	
		70
0.15	34	
		10
0.25	35	
		20
0.35	37	
		10
0.45	38	
		0
0.55	38	
		-50
0.65	33	

y x
 $\uparrow$ $\uparrow$
 $(x-t)$

Use $(x-t)$ graph to answer the following questions:

a) Determine one interval in which:

1- The velocity is increasing: at the interval [0-0.2]s

2- The velocity is decreasing: at the interval [0.6-0.7]s

3- The velocity is constant: at the interval [0.4-0.5]s

b) Find the instantaneous velocity at $t = 0.4$ s from the slope of the tangent of the $(x-t)$

graph: $v = \frac{\Delta x}{\Delta t} = \frac{15.6 - 7}{0.465 - 0.24} = 38.2 \text{ cm/s}$

Fill in table (4) below using data from table (1), and then answer the following questions:

Table (4)		«أخذ فترات متباينة»
Average velocity $\bar{v} = \frac{\Delta x}{\Delta t}$ (cm/s)		
$\bar{v}_{7-1} = \frac{x_7 - x_1}{t_7 - t_1} =$	35.83	
$\bar{v}_{6-2} = \frac{x_6 - x_2}{t_6 - t_2} =$	37	
$\bar{v}_{5-3} = \frac{x_5 - x_3}{t_5 - t_3} =$	37.5	

c) The midpoint for the given intervals is $t_{\text{mid}} = 0.4$ s.

d) As the time interval is shortened, is there any relation between average velocities in table (4) and instantaneous velocity at the midpoint? Yes ; the instantaneous velocity at the midpoint

approximately equals the average velocity for the smallest interval

e) Write down the approximate instantaneous velocity at t_{mid} from table (4).

* $v(t_{\text{mid}}) \cong \bar{v}$ at smallest interval

$$v_{t=0.4} \cong \bar{v}_{5-3} = 37.5 \text{ cm/s}$$

f) Compare this result with that you have found in question (b) above?

$$\text{p.e} = \left| \frac{\begin{matrix} (v_{\text{in Qb}}) \\ 38.2 \end{matrix} - \begin{matrix} (v_{\text{in Qc}}) \\ 37.5 \end{matrix}}{\begin{matrix} (38.2 + 37.5)/2 \\ (\bar{v}) \end{matrix}} \right| * 100\% = 1.85\%$$

Use $(v-t)$ graph to answer the following questions:

g) Determine one interval in which the acceleration is positive: [0.2-0.3] s


h) Find the instantaneous velocity at $t = 0.4$ sec from the graph. 37.4 cm/s

i) And compare it with the result in question (b).

$$\text{p.e} = \left| \frac{38.2 - 37.4}{(38.2 + 37.4)/2} \right| * 100\% = 2.12\%$$

j) Calculate the area under the instantaneous velocity in the interval [0.2-0.4] sec.

$$\text{Area} = \frac{(35 + 34.5)}{2} * 0.05 + \frac{(37 + 35)}{2} * 0.05 + \frac{(37 + 37.4)}{2} * 0.05 = 5.4 \text{ cm}$$



k) What does this area represent? Δx_{4-2} (the distance moved in the interval [0.2-0.4])

l) Compare it with the distance moved in the interval [0.2-0.4] sec from table (1).

$$\text{p.e} = \left| \frac{7.2 - 5.4}{(7.2 + 5.4)/2} \right| * 100\% = 28.57\%$$


Use Table (3) to answer the following questions:

m)


How does the acceleration change from one interval to the other (is it uniform or regular)?

The acceleration is changing irregularly.

Total distance (X) Versus Total time (t) graph



Average velocity ($\bar{v}$) versus (t) graph



Force and Motion: Newton's Second Law

Name: بشار عبد القادر العقاب

Partners: 10

Instructor: سماح النمر

ID#: 1440224

Section: 16

Purpose: To investigate Newton's second law: How a given force accelerates different masses and how different forces accelerate a given mass.


In figure (1)

M : mass of glider

m_a : added mass on glider

m_h : hanging mass

Figure 1



The theoretical equation of motion for the system in figure (1) is: $m_h g = (M + m_a + m_h) a$

Part (I): Acceleration and added mass with constant driving force.

Fill in table (1) with data from your experiment.

Table (1) m_h (hanging mass) = 50g

Glider's mass = 0.1 kg

Air pressure# (قوة الضغط)	Added mass m_a (kg)	Acceleration a (m/s ²)	$1/a$ (m/s ²) ⁻¹
4	0.000	3.189	0.3136
5	0.020	2.825	0.3539
6	0.050	2.206	0.4533
7	0.100	1.823	0.5485

(إضافة كتلة على عربة الغلدر)

Make a graph for m_a versus $1/a$. Then answer the following questions.

a) What is your conclusion about the way in which the acceleration depends on the magnitude of the added mass? When we increase the magnitude of the added mass the acceleration decreases. (the relation is inverse and linear)

b) Find the slope of your (m_a - $1/a$) graph. Slope = $\frac{\Delta m_a}{\Delta (1/a)} = \frac{0.0385 - 0.025}{0.4 - 0.37} = 0.45 \text{ N}$

What does the slope represent? The slope represents the driving force ($m_h g$). (وزن الكتلة المعلقة)

c) Determine the value of the glider mass (M) from the (m_a - $1/a$) graph. And compare it with the real value. * $y = \text{slope} x + y_{\text{intercept}}$ when $ma = 0$ $\Rightarrow 0 = 0.45 \times 0.3136 + y \Rightarrow y\text{-intercept} = -0.14 \text{ Kg}$

* Slope = $m_h g \Rightarrow m_h = 0.45 / 9.81 \Rightarrow m_h = 0.0459 \text{ Kg}$ * $|y\text{-intercept}| = M + m_h \Rightarrow M = 0.0941 \text{ Kg}$

* P.e = $\left| \frac{M_{\text{acc}} - M}{M_{\text{acc}}} \right| \times 100\% = \left| \frac{0.1 - 0.0941}{0.1} \right| \times 100\% = 5.9\%$

Part (II): Acceleration and driving force with constant total mass.

Fill in table (2) with the acceleration values from your experiment. Then, draw a graph for $m_h g$ versus a .

Table (2) M (mass of glider) = 100 g : ثبوت

Air pressure#	m_a (g)	m_h (g)	$m_h g$ (dyne)	a (cm/s ²)
7	100	50	49000	185.1
6	50	100	98000	360.1
5	20	130	127400	527.1
4	0	150	147000	608

a) What is your conclusion about the way in which the acceleration depends on the magnitude of the hanging mass?

When we increase the magnitude of the hanging mass, the acceleration increases (the relation is direct and linear).


b) Find the slope of your $m_h g$ versus a graph. What does the slope represent?

$$\text{Slope} = \frac{\Delta m_h g}{\Delta a} = \frac{(8.84 - 6.8) \times 10^4}{350 - 270} = 255 \text{ g}; \text{ It represents the total Mass } (M + m_a + m_h).$$


c) Do you expect that the curve should pass through the origin? Explain your answer.

Yes, I do; When there is no hanging mass, the acceleration will be zero $\begin{matrix} (0, 0) \\ \downarrow \quad \downarrow \\ a \quad m_h g \end{matrix}$

" m_a Versus $1/a$ graph"



« m_g versus a graph »



LAB REPORT FOR EXPERIMENT 8

Date: 30/3/2017

Name: إسماعيل عبد القادر الحفاني

Partner's Name: عثمان ناصر الحفاني

Registration No: 1440224

Registration No: 1438579

Physics Section: 16

Instructor's Name: د. سماح النور

PHYSICS LAB EXPERIMENT 8: SIMPLE HARMONIC MOTION :
THE SIMPLE PENDULUM

I. PURPOSE :

Study the relation between the period of oscillation (T) and
the length of the pendulum (L)

قياس الجاذبية للكواكب

II. DATA AND DATA ANALYSIS :

1. Compute the average of the period for each length of the pendulum and record them in Table 8.1.
2. Fill in the average periods $\bar{T}$ and the lengths L in table 8.2.

Table 8.1

	$L_1 = 30$ cm	$L_2 = 45$ cm	$L_3 = 60$ cm	$L_4 = 75$ cm	$L_5 = 90$ cm	$L_6 =$ cm
Trial No.	$t/20$ (s)	$t/20$ (s)	$t/20$ (s)	$t/20$ (s)	$t/20$ (s)	$t/20$ (s)
1	1.11	1.34	1.53	1.73	1.88	—
2	1.10	1.31	1.54	1.75	1.94	—
3	1.06	1.33	1.50	1.74	1.89	—
4	—	—	—	—	—	—
5	—	—	—	—	—	—
	$\bar{T}_1 = 1.09$ s	$\bar{T}_2 = 1.33$ s	$\bar{T}_3 = 1.52$ s	$\bar{T}_4 = 1.74$ s	$\bar{T}_5 = 1.90$ s	$\bar{T}_6 =$ — s

Table 8.2

L (cm)	$\bar{T}$ (s)	$\bar{T}^2$ (s ²)
30	1.09	1.1881
45	1.33	1.7689
60	1.52	2.3104
75	1.74	3.0276
90	1.90	3.61

3. Compute the square of the average period for each length and record it in Table 8.2.
4. Use your data in Table 8.1 to plot $\bar{T}$ versus L . What conclusion can you obtain from your graph?

The relation between $\bar{T}$ and L is direct but non-Linear.

5. Now plot $\bar{T}^2$ versus L using Table 8.2. What kind of relationship do you obtain?

The relation between $\bar{T}^2$ and L is direct and Linear.

6. Compute the slope of your graph plotted in 4 above.

$$\text{slope} = \frac{\Delta \bar{T}^2}{\Delta L} = \frac{2 - 1.625}{50 - 41} = 0.042 \text{ s}^2/\text{cm}$$


7. Using the value of the slope you obtained calculate g , the acceleration due to gravity.

$$\text{Slope} = \frac{4\pi^2}{g} = 0.042 \Rightarrow g = \frac{4\pi^2}{0.042} = 939.96 \text{ cm/s}^2$$


8. Estimate the error in g . $g_t = 981 \text{ cm/s}^2$

$$\text{p.e} = \left| \frac{981 - 939.96}{981} \right| * 100\% = 4.2\%$$

$\bar{T}$ vs L graph



T^2 v.s L graph.



LAB REPORT FOR EXPERIMENT 9

Date: 30/3/2017

Name: سليم عبد القادر الحفافي

Partner's Name: _____

Registration No: 1440224

Registration No: _____

Physics Section: 16

Instructor's Name: د. سعاد النور

PHYSICS LAB EXPERIMENT 9: THE LAWS OF GASES

I. PURPOSE :

Investigate the variation in the volume of entrapped gas (air) with pressure under constant room temperature, and find the

II. DATA AND DATA ANALYSIS : atmospheric pressure.

A. Boyle's Law

1. Enter the data in Table (9.1) below:

Table (9.1)

Scale Readings (mm)		L = B-X (mm)	h = Y-X (mm)	1/L (mm ⁻¹)
X	Y			
165	120	435	-45	2.30×10^{-3}
171	140	429	-31	2.33×10^{-3}
195	210	405	15	2.47×10^{-3}
211	288	389	77	2.57×10^{-3}
230	345	370	115	2.70×10^{-3}
245	400	355	155	2.82×10^{-3}
—	—	—	—	—
—	—	—	—	—
B = <u>600</u> mm				
Average Room Temp. = <u>20</u> °C				

هذا الخطأ لعدم إرفاق ورقة التقويم
التي تحتوي القراءات

2. Plot $1/L$ as independent variable against h . Use the graph to find the value of the atmospheric pressure $P_a \pm \Delta P_a$.

$P_a = 915 \text{ mm Hg}$ (y-intercept)

3. Calculate the pressure of the entrapped gas P in each case, using the relation: $P = (P_a + h)$ and also calculate the quantity PL and enter their values in Table (9.2) below:

Table (9.2)

L (mm)	$P = P_a + h$ (mm Hg)	PL (mmHg . mm)
435	$915 + (-45) = 870$	$378450 = 37.85 \times 10^4$
429	$915 + (-31) = 884$	$379236 = 37.92 \times 10^4$
405	$915 + 15 = 930$	$376650 = 37.67 \times 10^4$
389	$915 + 77 = 992$	$385888 = 38.59 \times 10^4$
370	$915 + 115 = 1030$	$381100 = 38.11 \times 10^4$
355	$915 + 155 = 1070$	$379850 = 37.99 \times 10^4$
—	—	—
—	—	—

4. Plot a second graph between L as independent variable and $P = (P_a + h)$. What do you conclude from such a curve?

$PV = \text{Constant}$
↑↑


The relation between (L) and (P) is an inverse and non-linear relation


5. Plot a third graph between L as independent variable against PL . What do you conclude from this graph?

$PV = \text{Constant}$
↑↑


The relation between (L) and (PL) is a constant relation,

h vs $(1/L)$ graph





(P) v.s (L) graph



(P_L) v.s (L) graph

LAB REPORT FOR EXPERIMENT 11

Date: 06/04/2017

Name: بشار عبدالمقار الحفالي

Partner's Name: _____

Registration No: 1440224

Registration No: _____

Physics Section: 16

Instructor's Name: أ.م.د. سعيد النور

PHYSICS LAB EXPERIMENT 11: SPECIFIC HEAT CAPACITY OF METALS

I. PURPOSE :

To Determine the specific heat capacity of unknown metal based on the principle of calorimetry.

II. DATA AND DATA ANALYSIS :

1. Enter the data in Table (11.1) below:

Table (11.1)

Specific heat capacity of calorimeter C_1	= 0.22	cal/g °C
Specific heat capacity of water C_w	= 1	cal/g °C
Mass of calorimeter M_1	= 63.15	g
Mass of calorimeter + water	= 96.96	g
Mass of water M_w	= 33.81	g
Temp. of calorimeter. +water T_1	= 21	°C
Temp. of metal T_2	= 95	°C
Final Equilibrium Temp. T_f	= 26	°C
Mass of metal M_2	= 36.66	g

2. What are the possible sources of error in this experiment?

1. error in measuring substance weight.

2. Loss of heat

3. Some personal errors.

3. Calculate the specific heat capacity (C_2) of the metal using the following equation:

Heat gained (by calorimeter + water) = Heat lost (by metal)

$$C_1 M_1 (T_f - T_1) + C_w M_w (T_f - T_w) = -C_2 M_2 (T_f - T_2)$$

$$(C_1 M_1 + C_w M_w) (T_f - T_1) = C_2 M_2 (T_2 - T_f)$$

$$(0.22 \times 63.15 + 1 \times 33.81) (26 - 21) = C_2 \times 36.66 (95 - 26)$$

$$\Rightarrow C_2 = 0.094 \text{ cal/g}^\circ\text{C}$$

4. Calculate ΔC_2 and express your final result as: $C_2 \pm \Delta C_2$. $\Delta M = 0.01 \text{ g}$

« derived quantity »

$\Delta T = 0.1^\circ\text{C}$

$$C_2 = \frac{(C_1 M_1 + C_w M_w) (T_f - T_1)}{M_2 (T_2 - T_f)} \quad \left(\frac{X Y}{M_2} \right)$$

$$X = C_1 M_1 + C_w M_w = 0.22 \times 63.15 + 1 \times 33.81 = 47.7 \text{ Cal/}^\circ\text{C}$$

$$\Delta X = \sqrt{(C_1 \Delta M_1)^2 + (C_w \Delta M_w)^2} = (0.22 \times 0.01)^2 + (1 \times 0.01)^2 = 0.01 \text{ Cal/}^\circ\text{C}$$

$$Y = T_f - T_1 = 26 - 21 = 5^\circ\text{C}$$

$$Z = T_2 - T_f = 95 - 26 = 69^\circ\text{C}$$

$$\Delta Y = \sqrt{(\Delta T_f)^2 + (\Delta T_1)^2} = 0.14^\circ\text{C}$$

$$\Delta Z = \sqrt{(\Delta T_2)^2 + (\Delta T_f)^2} = 0.14^\circ\text{C}$$

$$C_2 \pm \Delta C_2 = 0.094 \pm 0.0026 \text{ Cal/g}^\circ\text{C}$$

$$\Delta C_2 = \sqrt{\left(\frac{\Delta X}{X}\right)^2 + \left(\frac{\Delta Y}{Y}\right)^2 + \left(\frac{\Delta Z}{Z}\right)^2 + \left(\frac{\Delta M_2}{M_2}\right)^2} \times C_2$$

$$= \sqrt{\left(\frac{0.01}{47.7}\right)^2 + \left(\frac{0.14}{5}\right)^2 + \left(\frac{0.14}{69}\right)^2 + \left(\frac{0.01}{36.66}\right)^2} \times 0.094$$

5. What is the heat capacity of the metal sample?

$$C = C_2 M_2 = 0.094 \times 36.66 = 3.45 \text{ Cal/}^\circ\text{C}$$

6. How much heat is required to raise the temperature of 120g of Aluminum from 25 °C to 140 °F .

$$\begin{aligned}\Delta Q &= cM(T_f - T_i) \quad \Delta T & * T_f &= 1.8 T_c + 32 \\ &= 0.094 * 120 (60 - 25) & 140 &= 1.8 T_c + 32 \\ &= 394.8 \text{ Cal} & \Rightarrow T_c &= 60^\circ\text{C}\end{aligned}$$